Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Percivalle, E. et al., Viruses. 2020 Feb; 12(2): 157. | West Nile or Usutu Virus? A Three-Year Follow-Up of Humoral and Cellular Response in a Group of Asymptomatic Blood Donors | https://pubmed.ncbi.nlm.nih.gov/32013152/ |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Adotevi, O. et al., Clin Cancer Res.,2006; 12(10): 3158-67. | Immunogenic HLA-B*0702-restricted epitopes derived from human telomerase reverse transcriptase that elicit antitumor cytotoxic T-cell responses | https://www.ncbi.nlm.nih.gov/pubmed/16707616 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Alatrakchi N. et al., AIDS, 2002; 16(5): 713 - 717 | Strong CD4 Th1 responses to HIV and hepatitis C virus in HIV-infected long-term non-progressors co-infected with hepatitis C virus | https://www.ncbi.nlm.nih.gov/pubmed/14500478 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Almeida, J. R. et al.,Blood,2009; 113(25): 6351-6360. | Antigen sensitivity is a major determinant of CD8+ T-cell polyfunctionality and HIV-suppressive activity | https://www.ncbi.nlm.nih.gov/pubmed/19389882 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Almeida, J. R. et al., J. Exp. Med.,2007; 204(10): 2473-2485. | Superior control of HIV-1 replication by CD8+ T cells is reflected by their avidity, polyfunctionality, and clonal turnover | https://www.ncbi.nlm.nih.gov/pubmed/17893201 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Anguille, S. et al.,PLoS One,2012; 7(12): e51851 | Interleukin-15-induced CD56(+) myeloid dendritic cells combine potent tumor antigen presentation with direct tumoricidal potential. | https://www.ncbi.nlm.nih.gov/pubmed/23284789 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Ascough, S. et al., J Infect.,2014 ;68(2): 200-3 | Injectional anthrax infection due to heroin use induces strong immunological memory. | https://www.ncbi.nlm.nih.gov/pubmed/24513100 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Ascough, S.J. et al.,PLoS Pathog.,2014; 10(5): e1004085 | Anthrax lethal factor as an immune target in humans and transgenic mice and the impact of HLA polymorphism on CD4+ T cell immunity. | https://www.ncbi.nlm.nih.gov/pubmed/24788397 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Ayyoub M. et al., J. Immunol., 2002; 168(4):1717 - 1722 | Proteasome-assisted identification of a SSX-2-derived epitope recognized by tumor-reactive CTL infiltrating metastatic melanoma | https://www.ncbi.nlm.nih.gov/pubmed/11823502 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Ayyoub M. et al., J. Immunol., 2004; 172(11): 7206 - 7211 | Identification of an SSX-2 Epitope Presented by Dendritic Cells to Circulating Autologous CD4+ T Cells | https://www.ncbi.nlm.nih.gov/pubmed/15153546 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Bain C. et al., J. Virol., 2004; 78(19):10460 - 10469 | Memory T-Cell-Mediated Immune Responses Specific to an Alternative Core Protein in Hepatitis C Virus Infection | https://www.ncbi.nlm.nih.gov/pubmed/15367612 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Barragué, H. et al.,Medicine (Baltimore). 2017 Sep;96(39):e7915 | Chronic hepatitis E virus infection in a cirrhotic patient: A case report. | https://www.ncbi.nlm.nih.gov/pubmed/28953614 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Best, I. et al., Immunology,2009;128(1 Suppl):e777-86 | IFN-gamma production in response to Tax 161-233, and frequency of CD4+ Foxp3+ and Lin HLA-DRhigh CD123+ cells, discriminate HAM/TSP patients from asymptomatic HTLV-1-carriers in a Peruvian population. | https://www.ncbi.nlm.nih.gov/pubmed/19740339 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Beziaud, L. et al., Cancer Res.,2016 ; 76(14): 4100-4112 | Rapalogs Efficacy Relies on the Modulation of Antitumor T-cell Immunity | https://www.ncbi.nlm.nih.gov/pubmed/27197194 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Bolonaki, I. et al., J. Clin. Oncol.,2007; 25(19): 2727-2734. | Vaccination of Patients With Advanced Non-Small-Cell Lung Cancer With an Optimized Cryptic Human Telomerase Reverse Transcriptase Peptide | https://www.ncbi.nlm.nih.gov/pubmed/17602077 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Butt, N. M. et al., Haematologica,2005; 90(10): 1315-1323. | Circulating bcr-abl-specific CD8+ T cells in chronic myeloid leukemia patients and healthy subjects | https://www.ncbi.nlm.nih.gov/pubmed/16219567 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Calarota, S. A. et al., J. Immunol.,2008; 180(9): 5907-5915. | HIV-1-Specific T Cell Precursors with High Proliferative Capacity Correlate with Low Viremia and High CD4 Counts in Untreated Individuals | https://www.ncbi.nlm.nih.gov/pubmed/18424710 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Cassaniti,I. et al.,J Immunol Res. 2019 Feb 17;2019:4236503. | Evaluation of EBV- and HCMV-Specific T Cell Responses in Systemic Lupus Erythematosus (SLE) Patients Using a Normalized Enzyme-Linked Immunospot (ELISPOT) Assay. | https://www.ncbi.nlm.nih.gov/pubmed/30906789 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Calarota, S. A. et al.,Immunology,2013; 139(4): 533-44 | Detection of Epstein-Barr virus-specific memory CD4+ T cells using a peptide-based cultured enzyme-linked immunospot assay. | https://www.ncbi.nlm.nih.gov/pubmed/23560877 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Chu, K. K. et al.,Eur J Immunol.,2011 ; 41(1): 107-15 | CD4+ T-cell immunity to the Burkholderia pseudomallei ABC transporter LolC in melioidosis. | https://www.ncbi.nlm.nih.gov/pubmed/21182082 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Codecasa, L. et al., J Clin Microbiol.,2006; 44(6): 1944-50. | An in-house RD1-based enzyme-linked immunospot-gamma interferon assay instead of the tuberculin skin test for diagnosis of latent Mycobacterium tuberculosis infection | https://www.ncbi.nlm.nih.gov/pubmed/16757583 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Combadiere B. et al., J. Exp. Med., 2004; 199(11): 1585 - 1593 | Distinct time effects of vaccination on long-term proliferative and IFN-gamma-producing T cell memory to smallpox in humans | https://www.ncbi.nlm.nih.gov/pubmed/15184506 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Cools, N. et al., Mol Cancer,2006; 5: 49. | Sensitive detection of human papillomavirus type 16 E7-specific T cells by ELISPOT after multiple in vitro stimulations of CD8+ T cells with peptide-pulsed autologous dendritic cells | https://www.ncbi.nlm.nih.gov/pubmed/17067378 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | De Keersmaecker, B. et al., J. Leukoc. Biol.,2011 ; 89(6): 989-999 | The combination of 4-1BBL and CD40L strongly enhances the capacity of dendritic cells to stimulate HIV-specific T cell responses | https://www.ncbi.nlm.nih.gov/pubmed/21427207 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Decrion, A. Z. et al.,Immunology,2007; 121(3): 405-15 | A subset of functional effector-memory CD8+ T lymphocytes in human immunodeficiency virus-infected patients | https://www.ncbi.nlm.nih.gov/pubmed/17376193 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Dosset, M. et al., Clin Cancer Res., 2012; 18(22): 6284-95 | Universal cancer peptide-based therapeutic vaccine breaks tolerance against telomerase and eradicates established tumor | https://www.ncbi.nlm.nih.gov/pubmed/23032748 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Drillien R. et al., J. gen. Virol., 2004; 85(Pt 8): 2167 - 2175 | Modified vaccinia vifus Ankara induces moderate activation of human dendritic cells | https://www.ncbi.nlm.nih.gov/pubmed/15269355 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Durgeau, A. et al.,Nat Commun. 2018 Nov 30;9(1):5097. | Human preprocalcitonin self-antigen generates TAP-dependent and -independent epitopes triggering optimised T-cell responses toward immune-escaped tumours. | https://www.ncbi.nlm.nih.gov/pubmed/30504837 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Dufait, I. et al., Oncotarget,2015;6(14): 12369-82 | Ex vivo generation of myeloid-derived suppressor cells that model the tumor immunosuppressive environment in colorectal cancer. | https://www.ncbi.nlm.nih.gov/pubmed/25869209 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Elkord, E. et al., Int Immunol.,2005;17(10): 1315-25. | Differential CTLs specific for prostate-specific antigen in healthy donors and patients with prostate cancer | https://www.ncbi.nlm.nih.gov/pubmed/16141246 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Elkord, E. et al.,Immunology,2005; 114(2): 204-12 | Human monocyte isolation methods influence cytokine production from in vitro generated dendritic cells | https://www.ncbi.nlm.nih.gov/pubmed/15667565 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Farhi, D.  et al.,Arch Dermatol.,2009; 145(1): 38-45 | Non-Sexually Related Acute Genital Ulcers in 13 Pubertal Girls: A Clinical and Microbiological Study. | https://www.ncbi.nlm.nih.gov/pubmed/19153341 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Fostier,K. et al.,Oncotarget, 2018 Apr 17;9(29):20476-20489 | Impact of lenalidomide maintenance on the immune environment of multiple myeloma patients with low tumor burden after autologous stem cell transplantation. | https://www.ncbi.nlm.nih.gov/pubmed/29755666 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Galaine, J. et al., J. Immunol.,2016 ; 197(5): 1597-1608 | Heparan Sulfate Proteoglycans Promote Telomerase Internalization and MHC Class II Presentation on Dendritic Cells | https://www.ncbi.nlm.nih.gov/pubmed/27481844 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Gazagne A. et al., J. Immunol. Methods., 2003; 283(1-2): 91-98 | A Fluorospot assay to detect single T lymphocytes simultaneously producing multiple cytokines | https://www.ncbi.nlm.nih.gov/pubmed/14659902 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Godard B. et al., Hum. Immunol., 2004; 65(11): 1307-18 | Optimization of an elispot assay to detect cytomegalovirus-specific CD8+ T lymphocytes. | https://www.ncbi.nlm.nih.gov/pubmed/15556681 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Goovaerts, O. et al., PLoS One,2014 ; 9(11): e113101 | Antigen-specific interferon-gamma responses and innate cytokine balance in TB-IRIS. | https://www.ncbi.nlm.nih.gov/pubmed/25415590 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Grafmueller, S. et al., The Journal of Infectious Disease, 2012; 205: 1142 - 1146 | Differential Antigen Specificity of Hepatitis C Virus–Specific Interleukin 17– and Interferon {gamma}–Producing CD8+ T Cells During Chronic Infection | https://www.ncbi.nlm.nih.gov/pubmed/22351943 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Gupta, R. et al., Reprod Biol Endocrinol.,2009; 7: 38 | Host immune responses to chlamydial inclusion membrane proteins B and C in Chlamydia trachomatis infected women with or without fertility disorders | https://www.ncbi.nlm.nih.gov/pubmed/19397832 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Hamdi, H. et al., Arthritis Res Ther.,2006; 8(4): R114 | Inhibition of anti-tuberculosis T-lymphocyte function with tumour necrosis factor antagonists. | https://www.ncbi.nlm.nih.gov/pubmed/16859506 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Hoarau, J. J. et al.,PLoS One,2013; 8(12): e84695 | Identical strength of the T cell responses against E2, nsP1 and capsid CHIKV proteins in recovered and chronic patients after the epidemics of 2005-2006 in La Reunion Island. | https://www.ncbi.nlm.nih.gov/pubmed/24376836 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Hudak S. et al., J. Immuno., 2002; 169(3): 1189 - 1196 | Immune surveillance and effector functions of CCR10(+) skin homing T cells | https://www.ncbi.nlm.nih.gov/pubmed/12133939 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Ingram, R. J.et al., J. Immunol.,2010;184(7): 3814-3821 | Natural Exposure to Cutaneous Anthrax Gives Long-Lasting T Cell Immunity Encompassing Infection-Specific Epitopes | https://www.ncbi.nlm.nih.gov/pubmed/20208010 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Junwei, W. et al., Cancer Cell Int.,2016; 17: 10. | In vivo enhancement of the MAGE-specific cellular immune response by a recombinant MAGE1-MAGE3-TBHSP70 tumor vaccine | https://www.ncbi.nlm.nih.gov/pubmed/27330408 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Kalogerakou, F. et al.,Hippokratia,2008; 12(4): 230-5. | Detection of T cells secreting type 1 and type 2 cytokines in the peripheral blood of patients with oral lichen planus | https://www.ncbi.nlm.nih.gov/pubmed/19158967 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Kotsakis, A. et al., Ann. Onc., 2012; 23: 442 - 449 | Clinical outcome of patients with various advanced cancer types vaccinated with an optimized cryptic human telomerase reverse transcriptase (TERT) peptide: results of an expanded phase II study | https://www.ncbi.nlm.nih.gov/pubmed/21873272 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Laheurte, C. et al.,Oncoimmunology, 2016 May; 5(5): e1137416 | Immunoprevalence and magnitude of HLA-DP4 versus HLA-DR-restricted spontaneous CD4(+) Th1 responses against telomerase in cancer patients. | https://www.ncbi.nlm.nih.gov/pubmed/27467955 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Li, T. et al., PLoS One,2006; 1: e24. | Long-term persistence of robust antibody and cytotoxic T cell responses in recovered patients infected with SARS coronavirus. | https://www.ncbi.nlm.nih.gov/pubmed/17183651 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Lisziewicz, J. et al.,PLoS One,2012; 7(5): e35416 | Single DermaVir immunization: dose-dependent expansion of precursor/memory T cells against all HIV antigens in HIV-1 infected individuals. | https://www.ncbi.nlm.nih.gov/pubmed/22590502 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Lomas M. et al., Ann. Onc., 2004; 15(2): 324 - 329 | Phase I clinical trial of a human idiatypic p53 vaccine in patients with advanced malignancy | https://www.ncbi.nlm.nih.gov/pubmed/14760129 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Loyon, R. et al.,Front Immunol. 2019 Sep 6;10:2121. | Peripheral Innate Lymphoid Cells Are Increased in First Line Metastatic Colorectal Carcinoma Patients: A Negative Correlation With Th1 Immune Responses. | https://www.ncbi.nlm.nih.gov/pubmed/31555301 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Lu, S. Y. et al., World J Gastroenterol.,2004; 10(1): 53-7. | Superantigen-SEA gene modified tumor vaccine for hepatocellular carcinoma: an in vitro study. | https://www.ncbi.nlm.nih.gov/pubmed/14695768 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Lu, Z. Y. et al.,Exp Hematol.,2007 ; 35(3): 443-53 | B7-1 and 4-1BB ligand expression on a myeloma cell line makes it possible to expand autologous tumor-specific cytotoxic T cells in vitro. | https://www.ncbi.nlm.nih.gov/pubmed/17309825 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Mantegani, P. et al., Clin Med Res.,2006; 4(4): 266-72. | Comparison of an in-house and a commercial RD1-based ELISPOT-IFN-gamma assay for the diagnosis of Mycobacterium tuberculosis infection. | https://www.ncbi.nlm.nih.gov/pubmed/17210976 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Martinez, V. et al., BMC Infect Dis .,2007; 7: 83 | T-cell and serological responses to Erp, an exported Mycobacterium tuberculosis protein, in tuberculosis patients and healthy individuals. | https://www.ncbi.nlm.nih.gov/pubmed/17655752 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Montcuquet, N. et al., Immunology,2008;125(3): 320-30. | Regulatory T-cell expansion and function do not account for the impaired alloreactivity of ex vivo-expanded T cells | https://www.ncbi.nlm.nih.gov/pubmed/18445006 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Nicholas, R. S. et al., BMC Neurol.2015; 15: 72 | MS in South Asians in England: early disease onset and novel pattern of myelin autoimmunity. | https://www.ncbi.nlm.nih.gov/pubmed/25935418 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Onodi, F. et al.,Front Oncol. 2018 Nov 13;8:517. | High Therapeutic Efficacy of a New Survivin LSP-Cancer Vaccine Containing CD4+ and CD8+ T-Cell Epitopes. | https://www.ncbi.nlm.nih.gov/pubmed/30483475 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Pereira da Silva, T., et al., BMC Infect Dis.,2017;17:606 | Risk factors for increased immune reconstitution in response to Mycobacterium tuberculosis antigens in tuberculosis HIV-infected, antiretroviral-naïve patients | https://www.ncbi.nlm.nih.gov/pubmed/28874142 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Petanidis, S. et al., PLoS One,2013;8(9):e73616 | Differential expression of IL-17, 22 and 23 in the progression of colorectal cancer in patients with K-ras mutation: Ras signal inhibition and crosstalk with GM-CSF and IFN-gamma. | https://www.ncbi.nlm.nih.gov/pubmed/24040001 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Picard, E. et al.,Oncoimmunology. 2018 Oct 19;8(2):e1527498. | Circulating NKp46+ Natural Killer cells have a potential regulatory property and predict distinct survival in Non-Small Cell Lung Cancer. | https://www.ncbi.nlm.nih.gov/pubmed/30713781 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Pittet M. J. et al., J. Immunol., 2001; 166(12): 7634- 7640 | Ex vivo IFN-gamma secretion by circulating CD8 T lymphocytes: implications of a novel approach for T cell monitoring in infectious and malignant diseases | https://www.ncbi.nlm.nih.gov/pubmed/11390521 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Puissant-Lubrano, B. et al., J Clin Invest.,2010; 120(5): 1636-44. | Control of vaccinia virus skin lesions by long-term-maintained IFN-gamma+TNF-alpha+ effector/memory CD4+ lymphocytes in humans | https://www.ncbi.nlm.nih.gov/pubmed/20364089 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Purbhoo, M. A. et al., J Immunol., 2006; 176(12): 7308-16. | Quantifying and imaging NY-ESO-1/LAGE-1-derived epitopes on tumor cells using high affinity T cell receptors | https://www.ncbi.nlm.nih.gov/pubmed/16751374 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Purbhoo, M. A. et al., Mol. Cancer Ther.,2007; 6(7): 2081-2091. | The HLA A*0201-restricted hTERT540-548 peptide is not detected on tumor cells by a CTL clone or a high-affinity T-cell receptor | https://www.ncbi.nlm.nih.gov/pubmed/17620437 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Rinaldi, M. et al.,Thorax, 2012; 10.1136/thoraxjnl-2011-200690 | Antielastin B-cell and T-cell immunity in patients with chronic obstructive pulmonary disease | https://www.ncbi.nlm.nih.gov/pubmed/22442201 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Rozieres, A. et al., Allergy,2009 ; 64(4): 534-42 | Detection and quantification of drug-specific T cells in penicillin allergy | https://www.ncbi.nlm.nih.gov/pubmed/19154548 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Rubio-Godoy, V. et al., Proc Natl Acad Sci.,2001;98(18): 10302-7. | Discrepancy between ELISPOT IFN-gamma secretion and binding of A2/peptide multimers to TCR reveals interclonal dissociation of CTL effector function from TCR-peptide/MHC complexes half-life | https://www.ncbi.nlm.nih.gov/pubmed/11517329 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Saito, N. et al., Science Translational Medicine,2014; 6(245): 245ra95- | An annexin A1-FPR1 interaction contributes to necroptosis of keratinocytes in severe cutaneous adverse drug reactions. | https://www.ncbi.nlm.nih.gov/pubmed/25031270 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Samri, A. et al., Clin Vaccine Immunol.,2006; 13(6): 684-97. | Evaluation of the interlaboratory concordance in quantification of human immunodeficiency virus-specific T cells with a gamma interferon enzyme-linked immunospot assay | https://www.ncbi.nlm.nih.gov/pubmed/16760328 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Sauce D. et al., Blood, 2002; 99(4) : 1165 - 1173 | Retrovirus-mediated gene transfer in primary T lymphocytes impairs their anti-Epstein-Barr virus potential through both culture-dependent and selection process-dependent mechanisms | https://www.ncbi.nlm.nih.gov/pubmed/11830462 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Sauce D. et al., Blood, 2003; 102(4): 1241 - 1244 | Retrovirus-mediated gene transfer in polyclonal T cells results in lower apoptosis and enhanced ex vivo cell expansion of CMV-reactive CD8 T cells as compared with EBV-reactive CD8 T cells | https://www.ncbi.nlm.nih.gov/pubmed/12730101 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Schaubert, K. L. et al., J. Immunol.,2007; 178(12): 7756-7766. | Availability of a Diversely Avid CD8+ T Cell Repertoire Specific for the Subdominant HLA-A2-Restricted HIV-1 Gag p2419-27 Epitope | https://www.ncbi.nlm.nih.gov/pubmed/17548613 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Silva, B. C. et al. , Mem Inst Oswaldo Cruz,2014; 109(8): 999-1004 | Mycobacterium tuberculosis epitope-specific interferon-g production in healthy Brazilians reactive and non-reactive to tuberculin skin test. | https://www.ncbi.nlm.nih.gov/pubmed/25494469 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Spehner, L. et al., International journal of molecular sciences vol. 21,18 6838. 17 Sep. 2020, doi:10.3390/ijms21186838 | Anti-Telomerase CD4+ Th1 Immunity and Monocytic-Myeloid-Derived-Suppressor Cells Are Associated with Long-Term Efficacy Achieved by Docetaxel, Cisplatin, and 5-Fluorouracil (DCF) in Advanced Anal Squamous Cell Carcinoma: Translational Study of Epitopes-HPV01 and 02 Trials. | https://pubmed.ncbi.nlm.nih.gov/32957741 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Sun Y. et al., J. Immunol. Methods, 2003; 272(1-2): 23 - 34 | A systematic comparison of methods to measure HIV-1 specific CD8 T cells | https://www.ncbi.nlm.nih.gov/pubmed/12505709 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Teixeira, L. et al.,Clin Cancer Res. 2019 Sep 26. doi: 10.1158/1078-0432.CCR-19-1614. | A First-in-Human Phase I Study of INVAC-1, an Optimized Human Telomerase DNA Vaccine in Patients with Advanced Solid Tumors. | https://www.ncbi.nlm.nih.gov/pubmed/31558479 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Van Craenenbroeck, A. H. et al., Transplantation,2015;99(1):120-7 | Induction of cytomegalovirus-specific T cell responses in healthy volunteers and allogeneic stem cell recipients using vaccination with messenger RNA-transfected dendritic cells. | https://www.ncbi.nlm.nih.gov/pubmed/25050468 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Van Gulck, E. et al., PLoS One,2012; 7(5): e37792 | Immune and viral correlates of "secondary viral control" after treatment interruption in chronically HIV-1 infected patients. | https://www.ncbi.nlm.nih.gov/pubmed/22666392 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Van Gulck, E. et al.,Clin Dev Immunol., 2012: 184979 | Interleukin-12p70 expression by dendritic cells of HIV-1-infected patients fails to stimulate gag-specific immune responses. | https://www.ncbi.nlm.nih.gov/pubmed/22844321 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Van Gulck, E. R. et al., Blood,2006; 107(5): 1818-27. | Efficient stimulation of HIV-1-specific T cells using dendritic cells electroporated with mRNA encoding autologous HIV-1 Gag and Env proteins | https://www.ncbi.nlm.nih.gov/pubmed/16263796 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Van Gulck, E. R., et al, J. Virol,2008; 82(7): 3561-3573. | Efficient In Vitro Expansion of Human Immunodeficiency Virus (HIV)-Specific T-Cell Responses by gag mRNA-Electroporated Dendritic Cells from Treated and Untreated HIV Type 1-Infected Individuals | https://www.ncbi.nlm.nih.gov/pubmed/18234800 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Voelter, V.  et al., Int. Immunol.,2008; 20(8): 1087-1096. | Characterization of Melan-A reactive memory CD8+ T cells in a healthy donor. | https://www.ncbi.nlm.nih.gov/pubmed/18573812 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Vrecko, S. et al., Oncotarget. 2018 Oct 23;9(83): 35394-35407 | Personalized identification of tumor-associated immunogenic neoepitopes in hepatocellular carcinoma in complete remission after sorafenib treatment. | https://www.ncbi.nlm.nih.gov/pubmed/30459932 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Walton, S. M. et al., J Immunol.,2006;177(11): 8212-8. | Spontaneous CD8 T cell responses against the melanocyte differentiation antigen RAB38/NY-MEL-1 in melanoma patients | https://www.ncbi.nlm.nih.gov/pubmed/17114498 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Wei, J. et al., J Gen Virol.,2006; 87(Pt 11): 3393-6. | Identification of an HLA-A*0201-restricted cytotoxic T-lymphocyte epitope in rotavirus VP6 protein | https://www.ncbi.nlm.nih.gov/pubmed/17030875 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Weiss, L. et al.,PLoS One,2010; 5(7): e11659 | Relationship between regulatory T cells and immune activation in human immunodeficiency virus-infected patients interrupting antiretroviral therapy. | https://www.ncbi.nlm.nih.gov/pubmed/20657770 |
| Human IFNγ ELISpot Set | 856.051.001 / 005 / 010 / 015 / 020 | Zhang, Y. et al.,PLoS One,2015; 10(5): e0126075 | Dendritic-tumor fusion cells derived heat shock protein70-peptide complex has enhanced immunogenicity. | https://www.ncbi.nlm.nih.gov/pubmed/25961716 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | De Keersmaecker, B. et al., J Immunother Cancer. 2020; 8(1): e000329. | TriMix and tumor antigen mRNA electroporated dendritic cell vaccination plus ipilimumab: link between T-cell activation and clinical responses in advanced melanoma | https://pubmed.ncbi.nlm.nih.gov/32114500/ |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Zavattoni, M. et al., Microorganisms. 2020 Jan; 8(1): 56. | Zika Virus Infection in Pregnancy: Advanced Diagnostic Approaches in Dengue-Naive and Dengue-Experienced Pregnant Women and Possible Implication for Cross-Reactivity and Cross-Protection | https://pubmed.ncbi.nlm.nih.gov/31905661/ |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Adotevi, O. et al., Clin Cancer Res.,2006; 12(10): 3158-67. | Immunogenic HLA-B*0702-restricted epitopes derived from human telomerase reverse transcriptase that elicit antitumor cytotoxic T-cell responses | https://www.ncbi.nlm.nih.gov/pubmed/16707616 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Alatrakchi N. et al., AIDS, 2002; 16(5): 713 - 717 | Strong CD4 Th1 responses to HIV and hepatitis C virus in HIV-infected long-term non-progressors co-infected with hepatitis C virus | https://www.ncbi.nlm.nih.gov/pubmed/14500478 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Almeida, J. R. et al.,Blood,2009; 113(25): 6351-6360. | Antigen sensitivity is a major determinant of CD8+ T-cell polyfunctionality and HIV-suppressive activity | https://www.ncbi.nlm.nih.gov/pubmed/19389882 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Almeida, J. R. et al., J. Exp. Med.,2007; 204(10): 2473-2485. | Superior control of HIV-1 replication by CD8+ T cells is reflected by their avidity, polyfunctionality, and clonal turnover | https://www.ncbi.nlm.nih.gov/pubmed/17893201 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Anguille, S. et al.,PLoS One,2012; 7(12): e51851 | Interleukin-15-induced CD56(+) myeloid dendritic cells combine potent tumor antigen presentation with direct tumoricidal potential. | https://www.ncbi.nlm.nih.gov/pubmed/23284789 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Ascough, S. et al., J Infect.,2014 ;68(2): 200-3 | Injectional anthrax infection due to heroin use induces strong immunological memory. | https://www.ncbi.nlm.nih.gov/pubmed/24513100 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Ascough, S.J. et al.,PLoS Pathog.,2014; 10(5): e1004085 | Anthrax lethal factor as an immune target in humans and transgenic mice and the impact of HLA polymorphism on CD4+ T cell immunity. | https://www.ncbi.nlm.nih.gov/pubmed/24788397 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Ayyoub M. et al., J. Immunol., 2002; 168(4):1717 - 1722 | Proteasome-assisted identification of a SSX-2-derived epitope recognized by tumor-reactive CTL infiltrating metastatic melanoma | https://www.ncbi.nlm.nih.gov/pubmed/11823502 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Ayyoub M. et al., J. Immunol., 2004; 172(11): 7206 - 7211 | Identification of an SSX-2 Epitope Presented by Dendritic Cells to Circulating Autologous CD4+ T Cells | https://www.ncbi.nlm.nih.gov/pubmed/15153546 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Bain C. et al., J. Virol., 2004; 78(19):10460 - 10469 | Memory T-Cell-Mediated Immune Responses Specific to an Alternative Core Protein in Hepatitis C Virus Infection | https://www.ncbi.nlm.nih.gov/pubmed/15367612 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Best, I. et al., Immunology,2009;128(1 Suppl):e777-86 | IFN-gamma production in response to Tax 161-233, and frequency of CD4+ Foxp3+ and Lin HLA-DRhigh CD123+ cells, discriminate HAM/TSP patients from asymptomatic HTLV-1-carriers in a Peruvian population. | https://www.ncbi.nlm.nih.gov/pubmed/19740339 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Beziaud, L. et al., Cancer Res.,2016 ; 76(14): 4100-4112 | Rapalogs Efficacy Relies on the Modulation of Antitumor T-cell Immunity | https://www.ncbi.nlm.nih.gov/pubmed/27197194 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Bolonaki, I. et al., J. Clin. Oncol.,2007; 25(19): 2727-2734. | Vaccination of Patients With Advanced Non-Small-Cell Lung Cancer With an Optimized Cryptic Human Telomerase Reverse Transcriptase Peptide | https://www.ncbi.nlm.nih.gov/pubmed/17602077 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Butt, N. M. et al., Haematologica,2005; 90(10): 1315-1323. | Circulating bcr-abl-specific CD8+ T cells in chronic myeloid leukemia patients and healthy subjects | https://www.ncbi.nlm.nih.gov/pubmed/16219567 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Calarota, S. A. et al., J. Immunol.,2008; 180(9): 5907-5915. | HIV-1-Specific T Cell Precursors with High Proliferative Capacity Correlate with Low Viremia and High CD4 Counts in Untreated Individuals | https://www.ncbi.nlm.nih.gov/pubmed/18424710 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Calarota, S. A. et al.,Immunology,2013; 139(4): 533-44 | Detection of Epstein-Barr virus-specific memory CD4+ T cells using a peptide-based cultured enzyme-linked immunospot assay. | https://www.ncbi.nlm.nih.gov/pubmed/23560877 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Chu, K. K. et al.,Eur J Immunol.,2011 ; 41(1): 107-15 | CD4+ T-cell immunity to the Burkholderia pseudomallei ABC transporter LolC in melioidosis. | https://www.ncbi.nlm.nih.gov/pubmed/21182082 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Codecasa, L. et al., J Clin Microbiol.,2006; 44(6): 1944-50. | An in-house RD1-based enzyme-linked immunospot-gamma interferon assay instead of the tuberculin skin test for diagnosis of latent Mycobacterium tuberculosis infection | https://www.ncbi.nlm.nih.gov/pubmed/16757583 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Combadiere B. et al., J. Exp. Med., 2004; 199(11): 1585 - 1593 | Distinct time effects of vaccination on long-term proliferative and IFN-gamma-producing T cell memory to smallpox in humans | https://www.ncbi.nlm.nih.gov/pubmed/15184506 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Cools, N. et al., Mol Cancer,2006; 5: 49. | Sensitive detection of human papillomavirus type 16 E7-specific T cells by ELISPOT after multiple in vitro stimulations of CD8+ T cells with peptide-pulsed autologous dendritic cells | https://www.ncbi.nlm.nih.gov/pubmed/17067378 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | De Keersmaecker, B. et al., J. Leukoc. Biol.,2011 ; 89(6): 989-999 | The combination of 4-1BBL and CD40L strongly enhances the capacity of dendritic cells to stimulate HIV-specific T cell responses | https://www.ncbi.nlm.nih.gov/pubmed/21427207 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Decrion, A. Z. et al.,Immunology,2007; 121(3): 405-15 | A subset of functional effector-memory CD8+ T lymphocytes in human immunodeficiency virus-infected patients | https://www.ncbi.nlm.nih.gov/pubmed/17376193 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Drillien R. et al., J. gen. Virol., 2004; 85(Pt 8): 2167 - 2175 | Modified vaccinia vifus Ankara induces moderate activation of human dendritic cells | https://www.ncbi.nlm.nih.gov/pubmed/15269355 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Dufait, I. et al., Oncotarget,2015;6(14): 12369-82 | Ex vivo generation of myeloid-derived suppressor cells that model the tumor immunosuppressive environment in colorectal cancer. | https://www.ncbi.nlm.nih.gov/pubmed/25869209 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Elkord, E. et al., Int Immunol.,2005;17(10): 1315-25. | Differential CTLs specific for prostate-specific antigen in healthy donors and patients with prostate cancer | https://www.ncbi.nlm.nih.gov/pubmed/16141246 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Elkord, E. et al.,Immunology,2005; 114(2): 204-12 | Human monocyte isolation methods influence cytokine production from in vitro generated dendritic cells | https://www.ncbi.nlm.nih.gov/pubmed/15667565 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Farhi, D.  et al.,Arch Dermatol.,2009; 145(1): 38-45 | Non-Sexually Related Acute Genital Ulcers in 13 Pubertal Girls: A Clinical and Microbiological Study. | https://www.ncbi.nlm.nih.gov/pubmed/19153341 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Galaine, J. et al., J. Immunol.,2016 ; 197(5): 1597-1608 | Heparan Sulfate Proteoglycans Promote Telomerase Internalization and MHC Class II Presentation on Dendritic Cells | https://www.ncbi.nlm.nih.gov/pubmed/27481844 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Gazagne A. et al., J. Immunol. Methods., 2003; 283(1-2): 91-98 | A Fluorospot assay to detect single T lymphocytes simultaneously producing multiple cytokines | https://www.ncbi.nlm.nih.gov/pubmed/14659902 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Godard B. et al., Hum. Immunol., 2004; 65(11): 1307-18 | Optimization of an elispot assay to detect cytomegalovirus-specific CD8+ T lymphocytes. | https://www.ncbi.nlm.nih.gov/pubmed/15556681 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Goovaerts, O. et al., PLoS One,2014 ; 9(11): e113101 | Antigen-specific interferon-gamma responses and innate cytokine balance in TB-IRIS. | https://www.ncbi.nlm.nih.gov/pubmed/25415590 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Grafmueller, S. et al., The Journal of Infectious Disease, 2012; 205: 1142 - 1146 | Differential Antigen Specificity of Hepatitis C Virus–Specific Interleukin 17– and Interferon {gamma}–Producing CD8+ T Cells During Chronic Infection | https://www.ncbi.nlm.nih.gov/pubmed/22351943 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Gupta, R. et al., Reprod Biol Endocrinol.,2009; 7: 38 | Host immune responses to chlamydial inclusion membrane proteins B and C in Chlamydia trachomatis infected women with or without fertility disorders | https://www.ncbi.nlm.nih.gov/pubmed/19397832 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Hamdi, H. et al., Arthritis Res Ther.,2006; 8(4): R114 | Inhibition of anti-tuberculosis T-lymphocyte function with tumour necrosis factor antagonists. | https://www.ncbi.nlm.nih.gov/pubmed/16859506 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Hoarau, J. J. et al.,PLoS One,2013; 8(12): e84695 | Identical strength of the T cell responses against E2, nsP1 and capsid CHIKV proteins in recovered and chronic patients after the epidemics of 2005-2006 in La Reunion Island. | https://www.ncbi.nlm.nih.gov/pubmed/24376836 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Hudak S. et al., J. Immuno., 2002; 169(3): 1189 - 1196 | Immune surveillance and effector functions of CCR10(+) skin homing T cells | https://www.ncbi.nlm.nih.gov/pubmed/12133939 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Ingram, R. J.et al., J. Immunol.,2010;184(7): 3814-3821 | Natural Exposure to Cutaneous Anthrax Gives Long-Lasting T Cell Immunity Encompassing Infection-Specific Epitopes | https://www.ncbi.nlm.nih.gov/pubmed/20208010 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Kalogerakou, F. et al.,Hippokratia,2008; 12(4): 230-5. | Detection of T cells secreting type 1 and type 2 cytokines in the peripheral blood of patients with oral lichen planus | https://www.ncbi.nlm.nih.gov/pubmed/19158967 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Kotsakis, A. et al., Ann. Onc., 2012; 23: 442 - 449 | Clinical outcome of patients with various advanced cancer types vaccinated with an optimized cryptic human telomerase reverse transcriptase (TERT) peptide: results of an expanded phase II study | https://www.ncbi.nlm.nih.gov/pubmed/21873272 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Kroemer, M. et al., The Journal of infection vol. 82,2 (2021): 282-327. doi:10.1016/j.jinf.2020.08.036 | COVID-19 patients display distinct SARS-CoV-2 specific T-cell responses according to disease severity. | https://pubmed.ncbi.nlm.nih.gov/32853599 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Kroemer, M. et al., Oncoimmunology vol. 7,4 e1412030. 17 Jan. 2018, doi:10.1080/ 2162402X. 2017. 1412030 | SALL4 oncogene is an immunogenic antigen presented in various HLA-DR contexts. | https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5889287 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Laheurte, C. et al.,Oncoimmunology, 2016 May; 5(5): e1137416 | Immunoprevalence and magnitude of HLA-DP4 versus HLA-DR-restricted spontaneous CD4(+) Th1 responses against telomerase in cancer patients. | https://www.ncbi.nlm.nih.gov/pubmed/27467955 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Li, T. et al., PLoS One,2006; 1: e24. | Long-term persistence of robust antibody and cytotoxic T cell responses in recovered patients infected with SARS coronavirus. | https://www.ncbi.nlm.nih.gov/pubmed/17183651 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Lisziewicz, J. et al.,PLoS One,2012; 7(5): e35416 | Single DermaVir immunization: dose-dependent expansion of precursor/memory T cells against all HIV antigens in HIV-1 infected individuals. | https://www.ncbi.nlm.nih.gov/pubmed/22590502 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Lomas M. et al., Ann. Onc., 2004; 15(2): 324 - 329 | Phase I clinical trial of a human idiatypic p53 vaccine in patients with advanced malignancy | https://www.ncbi.nlm.nih.gov/pubmed/14760129 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Lu, S. Y. et al., World J Gastroenterol.,2004; 10(1): 53-7. | Superantigen-SEA gene modified tumor vaccine for hepatocellular carcinoma: an in vitro study. | https://www.ncbi.nlm.nih.gov/pubmed/14695768 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Lu, Z. Y. et al.,Exp Hematol.,2007 ; 35(3): 443-53 | B7-1 and 4-1BB ligand expression on a myeloma cell line makes it possible to expand autologous tumor-specific cytotoxic T cells in vitro. | https://www.ncbi.nlm.nih.gov/pubmed/17309825 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Mantegani, P. et al., Clin Med Res.,2006; 4(4): 266-72. | Comparison of an in-house and a commercial RD1-based ELISPOT-IFN-gamma assay for the diagnosis of Mycobacterium tuberculosis infection. | https://www.ncbi.nlm.nih.gov/pubmed/17210976 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Martinez, V. et al., BMC Infect Dis .,2007; 7: 83 | T-cell and serological responses to Erp, an exported Mycobacterium tuberculosis protein, in tuberculosis patients and healthy individuals. | https://www.ncbi.nlm.nih.gov/pubmed/17655752 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Montcuquet, N. et al., Immunology,2008;125(3): 320-30. | Regulatory T-cell expansion and function do not account for the impaired alloreactivity of ex vivo-expanded T cells | https://www.ncbi.nlm.nih.gov/pubmed/18445006 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Nicholas, R. S. et al., BMC Neurol.2015; 15: 72 | MS in South Asians in England: early disease onset and novel pattern of myelin autoimmunity. | https://www.ncbi.nlm.nih.gov/pubmed/25935418 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Petanidis, S. et al., PLoS One,2013;8(9):e73616 | Differential expression of IL-17, 22 and 23 in the progression of colorectal cancer in patients with K-ras mutation: Ras signal inhibition and crosstalk with GM-CSF and IFN-gamma. | https://www.ncbi.nlm.nih.gov/pubmed/24040001 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Pittet M. J. et al., J. Immunol., 2001; 166(12): 7634- 7640 | Ex vivo IFN-gamma secretion by circulating CD8 T lymphocytes: implications of a novel approach for T cell monitoring in infectious and malignant diseases | https://www.ncbi.nlm.nih.gov/pubmed/11390521 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Puissant-Lubrano, B. et al., J Clin Invest.,2010; 120(5): 1636-44. | Control of vaccinia virus skin lesions by long-term-maintained IFN-gamma+TNF-alpha+ effector/memory CD4+ lymphocytes in humans | https://www.ncbi.nlm.nih.gov/pubmed/20364089 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Purbhoo, M. A. et al., J Immunol., 2006; 176(12): 7308-16. | Quantifying and imaging NY-ESO-1/LAGE-1-derived epitopes on tumor cells using high affinity T cell receptors | https://www.ncbi.nlm.nih.gov/pubmed/16751374 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Purbhoo, M. A. et al., Mol. Cancer Ther.,2007; 6(7): 2081-2091. | The HLA A*0201-restricted hTERT540-548 peptide is not detected on tumor cells by a CTL clone or a high-affinity T-cell receptor | https://www.ncbi.nlm.nih.gov/pubmed/17620437 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Rinaldi, M. et al.,Thorax, 2012; 10.1136/thoraxjnl-2011-200690 | Antielastin B-cell and T-cell immunity in patients with chronic obstructive pulmonary disease | https://www.ncbi.nlm.nih.gov/pubmed/22442201 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Rozieres, A. et al., Allergy,2009 ; 64(4): 534-42 | Detection and quantification of drug-specific T cells in penicillin allergy | https://www.ncbi.nlm.nih.gov/pubmed/19154548 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Rubio-Godoy, V. et al., Proc Natl Acad Sci.,2001;98(18): 10302-7. | Discrepancy between ELISPOT IFN-gamma secretion and binding of A2/peptide multimers to TCR reveals interclonal dissociation of CTL effector function from TCR-peptide/MHC complexes half-life | https://www.ncbi.nlm.nih.gov/pubmed/11517329 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Saito, N. et al., Science Translational Medicine,2014; 6(245): 245ra95- | An annexin A1-FPR1 interaction contributes to necroptosis of keratinocytes in severe cutaneous adverse drug reactions. | https://www.ncbi.nlm.nih.gov/pubmed/25031270 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Samri, A. et al., Clin Vaccine Immunol.,2006; 13(6): 684-97. | Evaluation of the interlaboratory concordance in quantification of human immunodeficiency virus-specific T cells with a gamma interferon enzyme-linked immunospot assay | https://www.ncbi.nlm.nih.gov/pubmed/16760328 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Sauce D. et al., Blood, 2002; 99(4) : 1165 - 1173 | Retrovirus-mediated gene transfer in primary T lymphocytes impairs their anti-Epstein-Barr virus potential through both culture-dependent and selection process-dependent mechanisms | https://www.ncbi.nlm.nih.gov/pubmed/11830462 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Sauce D. et al., Blood, 2003; 102(4): 1241 - 1244 | Retrovirus-mediated gene transfer in polyclonal T cells results in lower apoptosis and enhanced ex vivo cell expansion of CMV-reactive CD8 T cells as compared with EBV-reactive CD8 T cells | https://www.ncbi.nlm.nih.gov/pubmed/12730101 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Schaubert, K. L. et al., J. Immunol.,2007; 178(12): 7756-7766. | Availability of a Diversely Avid CD8+ T Cell Repertoire Specific for the Subdominant HLA-A2-Restricted HIV-1 Gag p2419-27 Epitope | https://www.ncbi.nlm.nih.gov/pubmed/17548613 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Silva, B. C. et al. , Mem Inst Oswaldo Cruz,2014; 109(8): 999-1004 | Mycobacterium tuberculosis epitope-specific interferon-g production in healthy Brazilians reactive and non-reactive to tuberculin skin test. | https://www.ncbi.nlm.nih.gov/pubmed/25494469 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Sun Y. et al., J. Immunol. Methods, 2003; 272(1-2): 23 - 34 | A systematic comparison of methods to measure HIV-1 specific CD8 T cells | https://www.ncbi.nlm.nih.gov/pubmed/12505709 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Van Craenenbroeck, A. H. et al., Transplantation, 2015;99(1):120-7 | Induction of cytomegalovirus-specific T cell responses in healthy volunteers and allogeneic stem cell recipients using vaccination with messenger RNA-transfected dendritic cells. | https://www.ncbi.nlm.nih.gov/pubmed/25050468 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Van Gulck, E. et al., PLoS One,2012; 7(5): e37792 | Immune and viral correlates of "secondary viral control" after treatment interruption in chronically HIV-1 infected patients. | https://www.ncbi.nlm.nih.gov/pubmed/22666392 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Van Gulck, E. et al.,Clin Dev Immunol., 2012: 184979 | Interleukin-12p70 expression by dendritic cells of HIV-1-infected patients fails to stimulate gag-specific immune responses. | https://www.ncbi.nlm.nih.gov/pubmed/22844321 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Van Gulck, E. R. et al., Blood,2006; 107(5): 1818-27. | Efficient stimulation of HIV-1-specific T cells using dendritic cells electroporated with mRNA encoding autologous HIV-1 Gag and Env proteins | https://www.ncbi.nlm.nih.gov/pubmed/16263796 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Van Gulck, E. R., et al, J. Virol,2008; 82(7): 3561-3573. | Efficient In Vitro Expansion of Human Immunodeficiency Virus (HIV)-Specific T-Cell Responses by gag mRNA-Electroporated Dendritic Cells from Treated and Untreated HIV Type 1-Infected Individuals | https://www.ncbi.nlm.nih.gov/pubmed/18234800 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Voelter, V.  et al., Int. Immunol.,2008; 20(8): 1087-1096. | Characterization of Melan-A reactive memory CD8+ T cells in a healthy donor. | https://www.ncbi.nlm.nih.gov/pubmed/18573812 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Walton, S. M. et al., J Immunol.,2006;177(11): 8212-8. | Spontaneous CD8 T cell responses against the melanocyte differentiation antigen RAB38/NY-MEL-1 in melanoma patients | https://www.ncbi.nlm.nih.gov/pubmed/17114498 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Wei, J. et al., J Gen Virol.,2006; 87(Pt 11): 3393-6. | Identification of an HLA-A*0201-restricted cytotoxic T-lymphocyte epitope in rotavirus VP6 protein | https://www.ncbi.nlm.nih.gov/pubmed/17030875 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Weiss, L. et al.,PLoS One,2010; 5(7): e11659 | Relationship between regulatory T cells and immune activation in human immunodeficiency virus-infected patients interrupting antiretroviral therapy. | https://www.ncbi.nlm.nih.gov/pubmed/20657770 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Zhang, Y. et al.,PLoS One,2015; 10(5): e0126075 | Dendritic-tumor fusion cells derived heat shock protein70-peptide complex has enhanced immunogenicity. | https://www.ncbi.nlm.nih.gov/pubmed/25961716 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Garcia-Castillo, M. D. et al., J. Cell Sci.,2015; 128(13): 2373-2387 | Retrograde transport is not required for cytosolic translocation of the B-subunit of Shiga toxin | https://www.ncbi.nlm.nih.gov/pubmed/25977475 |
| Human IFNγ ELISpot Kit | 856.051.001PC/005PC | Junwei, W. et al., Cancer Cell Int.,2016; 17: 10. | In vivo enhancement of the MAGE-specific cellular immune response by a recombinant MAGE1-MAGE3-TBHSP70 tumor vaccine | https://www.ncbi.nlm.nih.gov/pubmed/27330408 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Trus, I. et al., Front Immunol. 2019; 10: 3077. | CpG-Recoding in Zika Virus Genome Causes Host-Age-Dependent Attenuation of Infection With Protection Against Lethal Heterologous Challenge in Mice | https://pubmed.ncbi.nlm.nih.gov/32038625/ |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Ishii, K. et al., Virology. 2006 Aug 1; 351(2): 368–380. | Induction of protective immunity against severe acute respiratory syndrome coronavirus (SARS-CoV) infection using highly attenuated recombinant vaccinia virus DIs | https://pubmed.ncbi.nlm.nih.gov/16678878/ |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Abadie, V. et al.,PLoS One,2009; 4(12): e8159 | Original encounter with antigen determines antigen-presenting cell imprinting of the quality of the immune response in mice | https://www.ncbi.nlm.nih.gov/pubmed/19997562 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Adotevi, O.et al., Blood,2010; 115(15): 3025-3032 | Targeting human telomerase reverse transcriptase with recombinant lentivector is highly effective to stimulate antitumor CD8 T-cell immunity in vivo | https://www.ncbi.nlm.nih.gov/pubmed/20130242 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Adriouch, S. et al., Front Microbiol.,2011; 2: 199 | Improved Immunological Tolerance Following Combination Therapy with CTLA-4/Ig and AAV-Mediated PD-L1/2 Muscle Gene Transfer. | https://www.ncbi.nlm.nih.gov/pubmed/22046170 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Ascough, S. et al., Front Microbiol.,2016;6:1506 | CD4+ T Cells Targeting Dominant and Cryptic Epitopes from Bacillus anthracis Lethal Factor | https://www.ncbi.nlm.nih.gov/pubmed/26779161 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Ascough, S. et al., PLoS Pathog.,2014; 10(5): e1004085 | Anthrax lethal factor as an immune target in humans and transgenic mice and the impact of HLA polymorphism on CD4+ T cell immunity. | https://www.ncbi.nlm.nih.gov/pubmed/24788397 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Attaf, M. et al., Sci Rep.,2016; 6: 35006. | alphabeta T cell receptor germline CDR regions moderate contact with MHC ligands and regulate peptide cross-reactivity. | https://www.ncbi.nlm.nih.gov/pubmed/27775030 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Bergwerf, I. et al.,BMC Biotechnol.,2009; 9: 1. | Reporter gene-expressing bone marrow-derived stromal cells are immune-tolerated following implantation in the central nervous system of syngeneic immunocompetent mice | https://www.ncbi.nlm.nih.gov/pubmed/19128466 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Bialkowski, L. et al., Sci Rep.,2016;6:22509 | Intralymphatic mRNA vaccine induces CD8 T-cell responses that inhibit the growth of mucosally located tumours | https://www.ncbi.nlm.nih.gov/pubmed/26931556 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Bonduelle, O. et al., J. Immunol., 2012; 188: 952 - 956 | Protective Effect of CX3CR1+ Dendritic Cells in a Vaccinia Virus Pulmonary Infection Model | https://www.ncbi.nlm.nih.gov/pubmed/22219332 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Calvet, C. Y. et al., Mol Ther Methods Clin Dev.,2014 ; 1: 14045 | Optimization of a gene electrotransfer procedure for efficient intradermal immunization with an hTERT-based DNA vaccine in mice | https://www.ncbi.nlm.nih.gov/pubmed/26015983 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Chen A. et al., J. Virol., 2005; 79(9): 5568 - 5576 | H-2 Kd-Restricted Hepatitis B Virus-Derived Epitope Whose Specific CD8+ T Lymphocytes Can Produce Gamma Interferon without Cytotoxicity | https://www.ncbi.nlm.nih.gov/pubmed/15827171 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Cox, F. et al., PLoS One,2015; 10(12): e0145243 | Protection against H5N1 Influenza Virus Induced by Matrix-M Adjuvanted Seasonal Virosomal Vaccine in Mice Requires Both Antibodies and T Cells. | https://www.ncbi.nlm.nih.gov/pubmed/26696245 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Dosset,M. et al.,Oncoimmunology. 2018 Mar 15;7(6):e1433981. | PD-1/PD-L1 pathway: an adaptive immune resistance mechanism to immunogenic chemotherapy in colorectal cancer. | https://www.ncbi.nlm.nih.gov/pubmed/29872568 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Dunachie, S.J. et al.,Sci Rep. 2017 Sep 22;7(1):12143. | Infection with Burkholderia pseudomallei - immune correlates of survival in acute melioidosis. | https://www.ncbi.nlm.nih.gov/pubmed/28939855 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Gaidot, A. et al., Blood,2011 ; 117(10): 2975-2983 | Immune reconstitution is preserved in hematopoietic stem cell transplantation coadministered with regulatory T cells for GVHD prevention | https://www.ncbi.nlm.nih.gov/pubmed/21193693 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Gallou, C. et al., Oncotarget,2016; 7(37): 59417-59428 | A general strategy to optimize immunogenicity of HLA-B*0702 restricted cryptic peptides from tumor associated antigens: Design of universal neo-antigen like tumor vaccines for HLA-B*0702 positive patients | https://www.ncbi.nlm.nih.gov/pubmed/27506946 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Gao, X. et al.,Mol Med Rep. 2018 Jul;18(1):315-321. | Gas-filled ultrasound microbubbles enhance the immunoactivity of the HSP70-MAGEA1 fusion protein against MAGEA1-expressing tumours. | https://www.ncbi.nlm.nih.gov/pubmed/29749485 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Guan, X. J. et al.,World J Gastroenterol.,2002; 8(2): 294-7 | Construction and characterization of an experimental ISCOMS-based hepatitis B polypeptide vaccine | https://www.ncbi.nlm.nih.gov/pubmed/11925610 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Hardet, R. et al., Mol Ther.,2015;24(1): 87-95 | Oral-tolerization Prevents Immune Responses and Improves Transgene Persistence Following Gene Transfer Mediated by Adeno-associated Viral Vector | https://www.ncbi.nlm.nih.gov/pubmed/26265250 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Herrmann, A. et al., J Clin Invest.,2014; 125(6): 2547 | CTLA4 aptamer delivers STAT3 siRNA to tumor-associated and malignant T cells. | https://www.ncbi.nlm.nih.gov/pubmed/26030229 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Herrmann, A. et al.,Cancer Res., 2010; 70 (19) :7455-7464. | Targeting Stat3 in the Myeloid Compartment Drastically Improves the In vivo Antitumor Functions of Adoptively Transferred T Cells | https://www.ncbi.nlm.nih.gov/pubmed/20841481 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Huang, J. et al.,BMC Immunol.,2012; 13: 50 | Residue analysis of a CTL epitope of SARS-CoV spike protein by IFN-gamma production and bioinformatics prediction. | https://www.ncbi.nlm.nih.gov/pubmed/22963340 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Johansen, P. et al., Clin. Vaccine Immunol.,2011; 18(6): 907-913 | Relief from Zmp1-Mediated Arrest of Phagosome Maturation Is Associated with Facilitated Presentation and Enhanced Immunogenicity of Mycobacterial Antigens | https://www.ncbi.nlm.nih.gov/pubmed/21471301 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Kujawski, M. et al., Cancer Res.,2010; 70(23): 9599-9610 | Targeting STAT3 in Adoptively Transferred T Cells Promotes Their In Vivo Expansion and Antitumor Effects | https://www.ncbi.nlm.nih.gov/pubmed/21118964 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Le Gouellec, A. et al.,Mol Ther.,2013; 21(5): 1076-86 | A safe bacterial microsyringe for in vivo antigen delivery and immunotherapy. | https://www.ncbi.nlm.nih.gov/pubmed/23531551 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Li H. et al., J. Immunol., 2005; 174(1): 195 - 204 | Generation of Murine CTL by a Hepatitis B Virus-Specific Peptide and Evaluation of the Adjuvant Effect of Heat Shock Protein Glycoprotein 96 and Its Terinal Fragments | https://www.ncbi.nlm.nih.gov/pubmed/15611241 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Li, H. T. et al., World J Gastroenterol.,2005; 11(19): 2858-63. | Enhancement of humoral immune responses to HBsAg by heat shock protein gp96 and its N-terminal fragment in mice | https://www.ncbi.nlm.nih.gov/pubmed/15902719 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Lin, T. et al., Clin Exp Immunol.,2006; 144(2): 319-25. | Enhanced immunogenicity and antitumour effects with heterologous prime-boost regime using vaccines based on MG7-Ag mimotope of gastric cancer. | https://www.ncbi.nlm.nih.gov/pubmed/16634806 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Luo, D. et al., Infect Immun.,2006; 74(5): 2734-41. | Protective immunity elicited by a divalent DNA vaccine encoding both the L7/L12 and Omp16 genes of Brucella abortus in BALB/c mice | https://www.ncbi.nlm.nih.gov/pubmed/16622210 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Mahnke, Y. D.et al.,Immunology,2005; 115(3): 325-36. | Maintenance of long-term tumour-specific T-cell memory by residual dormant tumour cells | https://www.ncbi.nlm.nih.gov/pubmed/15946250 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Martin Caballero, J. et al.,PLoS One,2012; 7(12): e52976 | Chimeric infectious bursal disease virus-like particles as potent vaccines for eradication of established HPV-16 E7-dependent tumors. | https://www.ncbi.nlm.nih.gov/pubmed/23300838 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | McIlroy, D. et al.,Mol Ther.,2009;17(8): 1473-81 | DNA/amphiphilic block copolymer nanospheres promote low-dose DNA vaccination. | https://www.ncbi.nlm.nih.gov/pubmed/19417740 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Musson, J. A. et al., Infect. Immun.,2010; (78): 4356-4362 | Repertoire of HLA-DR1-restricted CD4 T cell responses to capsular Caf1 antigen of Y. pestis HLA-transgenic mice | https://www.ncbi.nlm.nih.gov/pubmed/20660611 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Musson, J. A. et al.,J. Immunol.,2014 ; 193(12): 6041-6049 | CD4+ T Cell Epitopes of FliC Conserved between Strains of Burkholderia: Implications for Vaccines against Melioidosis and Cepacia Complex in Cystic Fibrosis | https://www.ncbi.nlm.nih.gov/pubmed/25392525 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Pere, H. et al., Blood,2011;118(18):4853-4862 | A CCR4 antagonist combined with vaccines induces antigen-specific CD8+ T cells and tumor immunity against self antigens | https://www.ncbi.nlm.nih.gov/pubmed/21908423 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Qiu, J. et al., Clin. Vaccine Immunol.,2011: CVI.00254-10 | Intranasal vaccination with recombinant Listeria {Delta}actA prfA* elicited robust systemic and pulmonary cellular responses, and secretory mucosal IgA | https://www.ncbi.nlm.nih.gov/pubmed/21270282 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Qiu, J. et al.,World J Gastroenterol.,2006; 12(3): 473-8. | Heat-shocked tumor cell lysate-pulsed dendritic cells induce effective anti-tumor immune response in vivo. | https://www.ncbi.nlm.nih.gov/pubmed/16489653 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Quigley, K. J. et al., Am J Respir Crit Care Med.,2015; 191(11): 1250-64 | Chronic Infection by Mucoid Pseudomonas aeruginosa Associated with Dysregulation in T-Cell Immunity to Outer Membrane Porin F. | https://www.ncbi.nlm.nih.gov/pubmed/25789411 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Rabu, C. et al., Oncoimmunology. 2019; 8(4): e1560919 | Cancer vaccines: designing artificial synthetic long peptides to improve presentation of class I and class II T cell epitopes by dendritic cells | https://www.ncbi.nlm.nih.gov/pubmed/30906653 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Ramakrishna L. et al., J. Virol., 2004; 78(17): 9174 - 9189 | Codon Optimization of the Tat Antigen of Human Immunodeficiency Virus Type 1 Generates Strong Immune Responses in Mice following Genetic Immunization | https://www.ncbi.nlm.nih.gov/pubmed/15308713 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Reynolds, C. et al., J. Immunol.2015 ; 194(10): 4814-4824 | T Cell Immunity to the Alkyl Hydroperoxide Reductase of Burkholderia pseudomallei: A Correlate of Disease Outcome in Acute Melioidosis | https://www.ncbi.nlm.nih.gov/pubmed/25862821 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Reynolds, C.J. et al.,Sci Rep., 2018 Jan 12;8(1):672 | T cell immunity to Zika virus targets immunodominant epitopes that show cross-reactivity with other Flaviviruses. | https://www.ncbi.nlm.nih.gov/pubmed/29330423 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Rojas, J. M. et al.,Vet Res.,2015 ; 45: 30 | Ovine and murine T cell epitopes from the non-structural protein 1 (NS1) of bluetongue virus serotype 8 (BTV-8) are shared among viral serotypes | https://www.ncbi.nlm.nih.gov/pubmed/24621015 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Rojas, J.M. et al.,Vet Res., 2017 Nov 21;48(1):79. | Vaccination with recombinant adenovirus expressing peste des petits ruminants virus-F or -H proteins elicits T cell responses to epitopes that arises during PPRV infection. | https://www.ncbi.nlm.nih.gov/pubmed/29157291 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Ru, Z. et al.,PLoS One,2012;7(3): e32247 | Development of a humanized HLA-A2.1/DP4 transgenic mouse model and the use of this model to map HLA-DP4-restricted epitopes of HBV envelope protein. | https://www.ncbi.nlm.nih.gov/pubmed/22403638 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Sandoval, F. et al., Science Translational Medicine,2013;5:172ra20 | Mucosal Imprinting of Vaccine-Induced CD8+ T Cells Is Crucial to Inhibit the Growth of Mucosal Tumors | https://www.ncbi.nlm.nih.gov/pubmed/23408053 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Seyed, N. et al.,PLoS One, 2014; 9(10): e108848 | Immunogenicity evaluation of a rationally designed polytope construct encoding HLA-A*0201 restricted epitopes derived from Leishmania major related proteins in HLA-A2/DR1 transgenic mice: steps toward polytope vaccine | https://www.ncbi.nlm.nih.gov/pubmed/25310094 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Shi, T. D. et al., World J Gastroenterol.,2004; 10(8): 1222-6. | Therapeutic polypeptides based on HBcAg(18-27) CTL epitope can induce antigen-specific CD(8)(+) CTL-mediated cytotoxicity in HLA-A2 transgenic mice | https://www.ncbi.nlm.nih.gov/pubmed/15069731 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Singh, M. et al.,PLoS One,2012; 7(6): e38491. | An improved protocol for efficient engraftment in NOD/LTSZ-SCIDIL-2Rgammanull mice allows HIV replication and development of anti-HIV immune responses. | https://www.ncbi.nlm.nih.gov/pubmed/22675567 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Someya K. et al., J. Virol., 2004; 78(18): 9842 - 9853 | A Consecutive Priming-Boosting Vaccination of Mice with Simian Immunodeficiency Virus (SIV) gag / pol DNA and Recombinant Vaccinia Virus Strain Dis Elicits Effective Anti-SIV Immunity | https://www.ncbi.nlm.nih.gov/pubmed/15331719 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Thalmensi, J. et al.,Oncoimmunology,2016; 5(3): e1083670. | Anticancer DNA vaccine based on human telomerase reverse transcriptase generates a strong and specific T cell immune response. | https://www.ncbi.nlm.nih.gov/pubmed/27141336 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Tian, G. et al.,World J Gastroenterol.,2004;10(2): 200-4 | Antitumor immunopreventive effect in mice induced by DNA vaccine encoding a fusion protein of alpha-fetoprotein and CTLA4. | https://www.ncbi.nlm.nih.gov/pubmed/14716822 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Tran, L. et al., J Biomed Biotechnol.,2012: 878657. | The immunogenicity of the tumor-associated antigen alpha-fetoprotein is enhanced by a fusion with a transmembrane domain | https://www.ncbi.nlm.nih.gov/pubmed/22500109 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Tran, T. et al., Clin. Cancer Res.,2016; 22(16): 4133-4144 | A Therapeutic Her2/neu Vaccine Targeting Dendritic Cells Preferentially Inhibits the Growth of Low Her2/neu-Expressing Tumor in HLA-A2 Transgenic Mice | https://www.ncbi.nlm.nih.gov/pubmed/27006496 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Tsunoda, I. et al., J Virol.,2005; 79(23): 14640-6. | Central nervous system pathology caused by autoreactive CD8+ T-cell clones following virus infection | https://www.ncbi.nlm.nih.gov/pubmed/16282464 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Waeckerle-Men, Y. et al., Nephrol Dial Transplant., 2007; 22: 1527 - 1536 | PD-L1 partially protects renal tubular epithelial cells from the attack of CD8+cytotoxic T cells | https://www.ncbi.nlm.nih.gov/pubmed/17339272 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Yan, L. et al., Infect. Immun.,2008; IAI.00245-08. | Selected prfA* mutations in recombinant attenuated Listeria monocytogenes augment expression of foreign immunogens and enhance vaccine-elicited humoral and cellular immune responses. | https://www.ncbi.nlm.nih.gov/pubmed/18474644 |
| Murine IFNγ ELISpot Set | 862.031.001 / 005 / 010 / 015 / 020 | Zhang, Y. et al., Oncol Rep.,2015; 33(6): 2695-702 | Enhanced antitumor immunity of nanoliposome-encapsulated heat shock protein 70 peptide complex derived from dendritic tumor fusion cells. | https://www.ncbi.nlm.nih.gov/pubmed/25963968 |
| Murine IFNγ ELISpot Kit | 862.031.001PC/005PC | Van Hoecke, L. et al., Mol Ther Nucleic Acids. 2020 Dec 4; 22: 373–381. | The Opposing Effect of Type I IFN on the T Cell Response by Non-modified mRNA-Lipoplex Vaccines Is Determined by the Route of Administration | https://pubmed.ncbi.nlm.nih.gov/33230442/ |
| Murine IFNγ ELISpot Kit | 862.031.001PC/005PC | Van Hoecke, L. et al., Oncoimmunology. 2020; 9(1): 1802968. | Recombinant viruses delivering the necroptosis mediator MLKL induce a potent antitumor immunity in mice | https://pubmed.ncbi.nlm.nih.gov/32923163/ |
| Murine IFNγ ELISpot Kit | 862.031.001PC/005PC | Huang, J. et al., Vaccine. 2006 Jun 5; 24(23): 4905–4913. | Immunization with SARS-CoV S DNA vaccine generates memory CD4+ and CD8+ T cell immune responses | https://pubmed.ncbi.nlm.nih.gov/16621188/ |
| Murine IFNγ ELISpot Kit | 862.031.001PC/005PC | Joe, P.T. et al., J Transl Med. 2019 Jul 25;17(1):242. | Intranodal administration of mRNA encoding nucleoprotein provides cross-strain immunity against influenza in mice. | https://www.ncbi.nlm.nih.gov/pubmed/31345237 |
| Murine IFNγ ELISpot Kit | 862.031.001PC/005PC | Musson, J. A. et al., Infect. Immun.,2010; (78): 4356-4362 | Repertoire of HLA-DR1-restricted CD4 T cell responses to capsular Caf1 antigen of Y. pestis HLA-transgenic mice | https://www.ncbi.nlm.nih.gov/pubmed/20660611 |
| Murine IFNγ ELISpot Kit | 862.031.001PC/005PC | Ramakrishna L. et al., J. Virol., 2004; 78(17): 9174 - 9189 | Codon Optimization of the Tat Antigen of Human Immunodeficiency Virus Type 1 Generates Strong Immune Responses in Mice following Genetic Immunization | https://www.ncbi.nlm.nih.gov/pubmed/15308713 |
| Murine IFNγ ELISpot Kit | 862.031.001PC/005PC | Someya K. et al., J. Virol., 2004; 78(18): 9842 - 9853 | A Consecutive Priming-Boosting Vaccination of Mice with Simian Immunodeficiency Virus (SIV) gag / pol DNA and Recombinant Vaccinia Virus Strain Dis Elicits Effective Anti-SIV Immunity | https://www.ncbi.nlm.nih.gov/pubmed/15331719 |
| Murine IFNγ ELISpot Kit | 862.031.001PC/005PC | Tsunoda, I. et al., J Virol.,2005; 79(23): 14640-6. | Central nervous system pathology caused by autoreactive CD8+ T-cell clones following virus infection | https://www.ncbi.nlm.nih.gov/pubmed/16282464 |
| Murine IFNγ ELISpot Kit | 862.031.001PC/005PC | Van Hoecke L et al., Nat Commun. 2018 Aug 24;9(1):3417 | Treatment with mRNA coding for the necroptosis mediator MLKL induces antitumor immunity directed against neo-epitopes. | https://www.ncbi.nlm.nih.gov/pubmed/30143632 |
| Murine IFNγ ELISpot Kit | 862.031.001PC/005PC | Yan, L. et al., Infect. Immun.,2008; IAI.00245-08. | Selected prfA* mutations in recombinant attenuated Listeria monocytogenes augment expression of foreign immunogens and enhance vaccine-elicited humoral and cellular immune responses. | https://www.ncbi.nlm.nih.gov/pubmed/18474644 |

Diaclone

# IFNγ ELISpot third-party citations

| Product | Catalogue Number | Citation | Title | Pubmed Link |
|---|---|---|---|---|
| Murine IFNγ ELISpot Kit | 862.031.001PC/005PC | Patil, H.P. et al.,Vaccines (Basel). 2017 Jul 27;5(3). pii: E19. | Adjuvantation of Pulmonary-Administered Influenza Vaccine with GPI-0100 Primarily Stimulates Antibody Production and Memory B Cell Proliferation. | https://www.ncbi.nlm.nih.gov/pubmed/28749414 |
| Murine IFNγ ELISpot Kit | 862.031.001PC/005PC | Pollard, C. et al., Mol Ther..,2013; 21(1): 251-9 | Type I IFN counteracts the induction of antigen-specific immune responses by lipid-based delivery of mRNA vaccines. | https://www.ncbi.nlm.nih.gov/pubmed/23011030 |
| Rat IFNγ ELISpot Set | 867.011.001 / 005 / 010 / 015 / 020 | Ginzkey, C. et al., Acta Neurochir (Wien),2013; 155(1): 51-8; discussion 59 | Incomplete tumour control following DNA vaccination against rat gliomas expressing a model antigen. | https://www.ncbi.nlm.nih.gov/pubmed/23132370 |